

5 CLAIMS

10/600446

1. A method for the treatment of protozoan infections comprising administering a quinazolinone compound in a pharmaceutically effective amount.

10

2. A method for the treatment of protozoan infections as recited in claim 1, wherein said compound is febrifugine/isofebrifugine extracted from *Dichroa febrifuga* Lour plant.

15

3. A method for the treatment of a protozoan infection as recited in claim 2, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

20

4. A method for the treatment of protozoan infections as recited in claim 3, wherein said protozoa is of the genus *Plasmodium*.

25


5. A method for the treatment of protozoan infections as recited in claim 4, wherein said protozoa is *Plasmodium falciparum*.

SV search

1, 2-11, 27-31
~~Excluded from 1-20, 55-59~~

5 6. A method for the treatment of protozoan infections as
recited in claim 4, wherein said protozoa is *Plasmodium berghei*.

7. A method for the treatment of protozoan infections
comprising administering a quinazolinone compound in a
10 pharmaceutical effective amount, said compound having the
formula:




15 8. A method for the treatment of protozoan infections as
recited in claim 7, wherein said administration is selected from
the group consisting of oral, subcutaneous, intramuscular, and
intraperitoneal.

20 9. A method for the treatment of protozoan infections as
recited in claim 8, wherein said protozoa is of the genus
Plasmodium.

5 10. A method for the treatment of protozoan infections as
recited in claim 9, wherein said protozoa is *Plasmodium*
falciparum.

11. A method for the treatment of protozoan infections as
10 recited in claim 9, wherein said protozoa is *Plasmodium berghei*.

12. A method for the treatment of ^ηprotozoan infections
comprising a quinazolinone compound in a pharmaceutical
effective amount, said compound having the formula:




13. A method for the treatment of protozoan infections as
recited in claim 12, wherein said administration is selected
from the group consisting of oral, subcutaneous, intramuscular,
20 and intraperitoneal.

14. A method for the treatment of protozoan infections as
recited in claim 13, wherein said protozoa is of the genus
Plasmodium.

15. A method for the treatment of protozoan infections as recited in claim 14, wherein said protozoa is *Plasmodium falciparum*.

10 16. A method for the treatment of protozoan infections as recited in claim 14, wherein said protozoa is *Plasmodium berghei*.

17. A method for the treatment of protozoan infections
15 comprising a quinazolinone compound in a pharmaceutical effective amount, said compound having the formula:




18. A method for the treatment of protozoan infections as recited in claim 17, wherein said administration is selected
20 from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

5 19. A method for the treatment of protozoan infections as
recited in claim 18, wherein said protozoa is of the genus
Plasmodium.

20 20. A method for the treatment of protozoan infections as
recited in claim 19, wherein said protozoa is *Plasmodium*
falciparum.

21. A method for the treatment of protozoan infections as
recited in claim 19, wherein said protozoa is *Plasmodium*
15 *berghei*.

22. A method for the treatment of protozoan infections
comprising a quinazolinone compound in a pharmaceutical
effective amount, said compound having the formula:



23. A method for the treatment of protozoan infections as
recited in claim 22, wherein said administration is selected
from the group consisting of oral, subcutaneous, intramuscular,
and intraperitoneal.


25

5 24. A method for the treatment of protozoan infections as recited in claim 23, wherein said protozoa is of the genus *Plasmodium*.

25. A method for the treatment of protozoan infections as
10 recited in claim 24, wherein said protozoa is *Plasmodium falciparum*.

26. A method for the treatment of protozoan infections as recited in claim 24, wherein said protozoa is *Plasmodium*
15 *berghei*.

27. A method for the treatment of protozoan infections comprising a quinazolinone compound in a pharmaceutical effective amount, said compound having the formula:




28. A method for the treatment of protozoan infections as recited in claim 27, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

5 29. A method for the treatment of protozoan infections as
recited in claim 28, wherein said protozoa is of the genus
Plasmodium.

30. A method for the treatment of protozoan infections as
10 recited in claim 29, wherein said protozoa is *Plasmodium*
falciparum.

31. A method for the treatment of protozoan infections as
recited in claim 29, wherein said protozoa is *Plasmodium*
15 *berghei*.

32. A method for the treatment of protozoan infections
comprising a quinazolinone compound in a pharmaceutical
effective amount, said compound having the formula:



33. A method for the treatment of protozoan infections as
recited in claim 32, wherein said administration is selected

5 from the group consisting of oral, subcutaneous, intramuscular,
and intraperitoneal.

34. A method for the treatment of protozoan infections as
recited in claim 33, wherein said protozoa is of the genus
10 *Plasmodium*.


35. A method for the treatment of protozoan infections as
recited in claim 34, wherein said protozoa is *Plasmodium*
falciparum.

15

36. A method for the treatment of protozoan infections as
recited in claim 34, wherein said protozoa is *Plasmodium*
berghei.

20

37. A method for the treatment of protozoan infections
comprising a quinazolinone compound in a pharmaceutical
effective amount, said compound having the formula:



25


38. A method for the treatment of protozoan infections as recited in claim 37, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

39. A method for the treatment of protozoan infections as recited in claim 38, wherein said protozoa is of the genus *Plasmodium*.

40. A method for the treatment of protozoan infections as recited in claim 39, wherein said protozoa is *Plasmodium falciparum*.

41. A method for the treatment of protozoan infections as recited in claim 39, wherein said protozoa is *Plasmodium berghei*.

42. A method for the treatment of protozoan infections comprising a quinazolinone compound in a pharmaceutical effective amount, said compound having the formula:




43. A method for the treatment of protozoan infections as recited in claim 42, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

44. A method for the treatment of protozoan infections as recited in claim 43, wherein said protozoa is of the genus *Plasmodium*.

45. A method for the treatment of protozoan infections as recited in claim 44, wherein said protozoa is *Plasmodium falciparum*.

46. A method for the treatment of protozoan infections as recited in claim 44, wherein said protozoa is *Plasmodium berghei*.

47. A method for the treatment of protozoan infections comprising a quinazolinone compound in a pharmaceutical effective amount, said compound having the formula:



48. A method for the treatment of protozoan infections as recited in claim 47, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

10

49. A method for the treatment of protozoan infections as recited in claim 48, wherein said protozoa is of the genus *Plasmodium*.


15

50. A method for the treatment of protozoan infections as recited in claim 49, wherein said protozoa is *Plasmodium falciparum*.

20

51. A method for the treatment of protozoan infections as recited in claim 50, wherein said protozoa is *Plasmodium berghei*.

5 52. A method for the treatment of protozoan infections comprising a quinazolinone compound in a pharmaceutical effective amount, said compound having the formula:



10 53. A method for the treatment of protozoan infections as recited in claim 52, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.


15 54. A method for the treatment of protozoan infections as recited in claim 53, wherein said protozoa is of the genus *Plasmodium*.

20 55. A method for the treatment of protozoan infections as recited in claim 54, wherein said protozoa is *Plasmodium falciparum*.

5 56. A method for the treatment of protozoan infections as recited in claim 54, wherein said protozoa is *Plasmodium berghei*.

5

57. A method for the treatment of protozoan infections comprising a quinazolinone compound in a pharmaceutical effective amount, said compound having the formula:



10

58. A method for the treatment of protozoan infections as recited in claim 57, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

15


59. A method for the treatment of protozoan infections as recited in claim 58, wherein said protozoa is of the genus *Plasmodium*.

20

60. A method for the treatment of protozoan infections as recited in claim 59, wherein said protozoa is *Plasmodium falciparum*.

5 61. A method for the treatment of protozoan infections as recited in claim 59, wherein said protozoa is *Plasmodium berghei*.

62. A method for the treatment of protozoan infections
10 comprising a quinazolinone compound in a pharmaceutical effective amount, said compound having the formula:




15 63. A method for the treatment of protozoan infections as recited in claim 62, wherein said administration is selected from the group consisting of oral, subcutaneous, intramuscular, and intraperitoneal.

20 64. A method for the treatment of protozoan infections as recited in claim 63, wherein said protozoa is of the genus *Plasmodium*.

5 65. A method for the treatment of protozoan infections as
recited in claim 64, wherein said protozoa is *Plasmodium*
falciparum.

66. A method for the treatment of protozoan infections as
10 recited in claim 64, wherein said protozoa is *Plasmodium*
berghei.

67. A method for the treatment of protozoan infections
comprising a quinazolinone compound in a pharmaceutical
15 effective amount, said compound having the formula:





68. A method for the treatment of protozoan infections as
recited in claim 67, wherein said administration is selected
.0 from the group consisting of oral, subcutaneous, intramuscular,
and intraperitoneal.


5 69. A method for the treatment of protozoan infections as
recited in claim 68, wherein said protozoa is of the genus
Plasmodium.

70. A method for the treatment of protozoan infections as
10 recited in claim 69, wherein said protozoa is *Plasmodium*
falciparum.



71. A method for the treatment of protozoan infections as
recited in claim 69, wherein said protozoa is *Plasmodium*
15 *berghei*.





7





12



17



22



27

